



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 314 662 B1

(12)

EUROPEAN PATENT SPECIFICATION

(49) Date of publication of patent specification: 19.11.92 (51) Int. Cl.5: C07F 9/38

(21) Application number: 88870162.0

(22) Date of filing: 25.10.88

(54) Process for producing N-phosphonomethylglycine.

(30) Priority: 26.10.87 US 112594

(43) Date of publication of application:
03.05.89 Bulletin 89/18

(45) Publication of the grant of the patent:
19.11.92 Bulletin 92/47

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(56) References cited:
DE-A- 2 366 060

CHEMICAL ABSTRACTS, vol. 101, no. 13, 24th
September 1984, page 681, abstract no.
111201q, Columbus, Ohio, US

(73) Proprietor: Monsanto Company
Patent Department 800 North Lindbergh Bou-
levard
St. Louis Missouri 63167-7020(US)

(72) Inventor: Riley, Dennis Patrick
15519 Country Ridge Drive
Chesterfield Missouri 63017(US)
Inventor: Rivers, Willie James, Jr.
1239 Waldron
University City Missouri 63130(US)

(74) Representative: Ernst, Hubert et al
Monsanto Services International S.A., Patent
Department, Avenue de Tervuren 270-272,
Letter Box No. 21
B-1150 Brussels(BE)

EP 0 314 662 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

This invention relates to a process for producing N-phosphonomethylglycine by the oxidation of N-phosphonomethyliminodiacetic acid using transition metal catalysts. More particularly, this invention relates to a reaction using molecular oxygen and a transition metal salt catalyst.

It is known in the art that N-phosphonomethylglycine can be produced by oxidizing N-phosphonomethyliminodiacetic acid using various oxidizing methods. U.S. patent 3,950,402 discloses a method wherein N-phosphonomethyliminodiacetic acid is oxidized to N-phosphonomethylglycine in aqueous media using a free oxygen-containing gas and a heterogeneous noble metal-based catalyst such as palladium, platinum or rhodium. U. S. patent 3,954,848 discloses the oxidation of N-phosphonomethyliminodiacetic acid with hydrogen peroxide and an acid such as sulfuric or acetic acid. U.S. patent 3,969,398 discloses the oxidation of N-phosphonomethyliminodiacetic acid using molecular oxygen and a heterogeneous activated carbon catalyst. Hungarian Patent Application No. 011706 discloses the oxidation of N-phosphonomethyliminodiacetic acid with peroxide in the presence of metals or metal compounds.

R. J. Motekaitis, A. E. Martell, D. Hayes and W. W. Frenier, Can. J. Chem., 58, 1999 (1980) disclose the iron(III) or copper(II) catalysed oxidative dealkylation of ethylene diaminetetracetic acid (EDTA) and nitrilotriacetic acid (NTA), both of which have iminodiacetic acid groups. R. J. Motekaitis, X. B. Cox, III, P. Taylor, A. E. Martell, B. Miles and T. J. Tvedt, Can. J. Chem., 60, 1207 (1982) disclose that certain metal ions, such as Ca(II), Mg(II), Fe(II), Zn(II) and Ni(II) chelate with EDTA and stabilize against oxidation, thereby reducing the rate of oxidative dealkylation.

The present invention involves a process for the production of N-phosphonomethylglycine comprising contacting N-phosphonomethyliminodiacetic acid with a molecular oxygen-containing gas in the presence of a transition metal catalyst.

The process of this invention involves contacting N-phosphonomethyliminodiacetic acid with a transition metal catalyst in a mixture or solution. This mixture or solution is contacted with a molecular oxygen-containing gas while heating the reaction mass to a temperature sufficiently elevated to initiate and sustain the oxidation reaction of N-phosphonomethyliminodiacetic acid to produce N-phosphonomethylglycine.

The transition metal catalyst of the present invention can be any one or more of several transition metal compounds such as manganese, cobalt, iron, nickel, chromium, ruthenium, aluminum, molybdenum, vanadium and cerium. The catalysts can be in the form of salts such as manganese salts, e.g., manganese acetate, manganese sulfate; complexes such as manganese(II)bis(acetylacetonate) (Mn(II)(acac)_2); cobalt salts such as $\text{Co(II)(SO}_4\text{)}$, $\text{Co(II)(acetylacetonate)}$, CoCl_2 , CoBr_2 , $\text{Co(NO}_3\text{)}_2$ and cobalt acetate; cerium salts such as $(\text{NH}_4)_4\text{Ce(SO}_4\text{)}$ and $(\text{NH}_4)_2\text{Ce(NO}_3\text{)}_6$, iron salts such as $(\text{NH}_4)_2\text{Fe(SO}_4\text{)}_2$, iron(III)(dicyano)-bis(phenanthroline)₂-(tetrafluoro)borate salt and $\text{K}_3\text{Fe(CN)}_6$, and other metal salts such as NiBr_2 , CrCl_3 , $\text{RuCl}_2(\text{Me}_2\text{SO})$, RuBr_3 , $\text{Al(NO}_3\text{)}_3$, $\text{K}_4\text{Mo(CN)}_8$, $\text{VO(acetylacetonate)}_2$ and VOSO_4 . The catalyst can be added to the N-phosphonomethyliminodiacetic acid in the salt form, or a salt may be generated in situ by the addition of a source of a transition metal ion such as MnO_2 which dissolves in the reaction medium. The Mn(III)chloro(phthalocyaninato), however, is not catalytic, possibly because the phthalocyanine ligand covalently bonds to the Mn(III) and therefore inhibits the formation of N-phosphonomethyliminodiacetic acid/manganese complex in solution.

Manganese salts such as Mn(II), Mn(III) or Mn(IV) salts can be used individually, however, the reaction displays a delayed reaction initiation time (initiation period), e.g., there is a delay before any N-phosphonomethylglycine is produced. When a mixture of Mn(II) and Mn(III) salts are used as a catalyst system, the initiation is diminished or eliminated. A preferred manganese salt catalyst is a mixture of Mn(II) and Mn(III) salts in the range of 1:10 to 10:1 mole ratio of the Mn ions. A most preferred manganese catalyst salt is a 1:1 mole ratio of Mn(II) and Mn(III) ions in the form of manganese acetate salts. A preferred cobalt catalyst is a Co(II) salt such as $\text{Co(II)(SO}_4\text{)}$, Co(II)Cl_2 , Co(II)Br_2 , Co(II)(OH)_2 and Co(II)acetate.

The concentration of the transition metal catalyst in the reaction solution can vary widely, in the range of 0.1M to 0.0001M total metal ion concentration. For manganese, the reaction appears to have a first order dependency on the catalyst concentration, e.g., the reaction rate increases linearly as the catalyst concentration increases. The preferred concentration is in the range of about 0.01M to about 0.001M, which gives a suitably fast rate of reaction that can be easily controlled and favors selectivity to N-phosphonomethylglycine.

The reaction temperature is sufficient to initiate and sustain the oxidation reaction, in the range of about 25°C to 150°C. In general, as the reaction temperature increases, the reaction rate increases. To achieve an easily controlled reaction rate and favor selectivity to N-phosphonomethylglycine, a preferred temperature range is about 50°C to 120°C and a most preferred is in the range of about 70°C to 100°C. If a

temperature of above about 100° C is used, pressure will have to be maintained on the system to maintain a liquid phase.

The pressure at which this process is conducted can vary over a wide range. The range can vary from about atmospheric (101 kPa) to about 3000 psig (20700 kPa). A preferred range is about 30 psig (200 kPa) to about 1000 psig (about 6900 kPa). A most preferred range is from about 150 psig (about 1000 kPa) to 600 psig (about 4140 kPa).

The oxygen concentration, as designated by the partial pressure of oxygen (PO_2), in the reaction affects the reaction rate and the selectivity to the desired product, N-phosphonomethylglycine. As the PO_2 increases, the reaction rate generally increases and the selectivity to N-phosphonomethylglycine increases. The PO_2 can be increased by increasing the overall reaction pressure, or by increasing the molecular oxygen concentration in the molecular oxygen-containing gas. The PO_2 can vary widely, in the range of from 1 psig (6.9 kPa) to 3000 psig (20700 kPa). A preferred range is from 30 psig (207 kPa) to 1000 psig (6900 kPa).

The term "molecular oxygen-containing gas" means molecular oxygen gas or any gaseous mixture containing molecular oxygen with one or more diluents which are non-reactive with the oxygen or with the reactant or product under the conditions of reaction. Examples of such diluent gases are air, helium, argon, nitrogen, or other inert gas, or oxygen-hydrocarbon mixtures. A preferred molecular oxygen is undiluted oxygen gas.

The manner in which the solution or mixture of the N-phosphonomethyliminodiacetic acid is contacted with molecular oxygen can vary greatly. For example, the N-phosphonomethyliminodiacetic acid solution or mixture can be placed in a closed container with some free space containing molecular oxygen and shaken vigorously or agitated by stirring. Alternatively, the molecular oxygen can be continuously bubbled through the solution or mixture containing the transition metal catalyst using a straight tube or a tube with a fritted diffuser attached to it. The process of this invention only requires actively contacting the molecular oxygen-containing gas with the aqueous solution or mixture of the N-phosphonomethyliminodiacetic acid containing a transition metal catalyst.

The initial pH (pHi) of the reaction affects the reaction rate and the selectivity to N-phosphonomethylglycine. For example, with manganese, as the initial pH increases, the reaction rate increases, but the selectivity to N-phosphonomethylglycine decreases. The pHi of the reaction can vary widely, in the range of about 0.1 to about 7. A preferred range is about 1 to about 3 with manganese and about 0.1 to 3 with cobalt. A most preferred pH is the unadjusted pH of N-phosphonomethyliminodiacetic acid in a water solution which varies with the N-phosphonomethyliminodiacetic acid concentration and the reaction temperature.

The oxidation reaction can take place in a solution or slurry. For a solution, the initial concentration of the N-phosphonomethyliminodiacetic acid in the reaction mass is a function of the solubility of the N-phosphonomethyliminodiacetic acid in the solvent at both the desired reaction temperature and the pHi of the solution. As the solvent temperature and pH changes, the solubility of the N-phosphonomethyliminodiacetic acid changes. A preferred initial concentration of the N-phosphonomethyliminodiacetic acid is a saturated slurry containing a solvent system at reaction conditions, which maximize the yield of N-phosphonomethylglycine in the reaction mass. A preferred concentration of N-phosphonomethyliminodiacetic acid is in the range of about 1 to 50 wt.%. It is, of course, possible to employ very dilute solutions of N-phosphonomethyliminodiacetic acid, or slurries and mixtures.

The reaction is typically carried out in an aqueous solvent. The term aqueous solvent means solutions containing at least about 50 weight % water. The preferred aqueous solvent is distilled, deionized water.

The following examples are for illustration purposes only and are not intended to limit the scope of the claimed invention.

EXAMPLES

A series of runs were made to oxidize N-phosphonomethyliminodiacetic acid to N-phosphonomethylglycine. The reactions were conducted in a modified Fisher-Porter glass pressure apparatus or an Engineer Autoclave 300 ml pressure reactor in which a stirrer was installed in the head, as were three additional valved ports that were used as a sample port, a gas inlet, and a purged gas outlet. The stirrer maintained sufficient agitation to afford thorough gas-liquid mixing. The temperature was controlled by immersing the reactor in a constant temperature oil bath. The indicated amount of transition metal catalyst was dissolved or suspended in a distilled, deionized water solution containing the indicated amount of N-phosphonomethyliminodiacetic acid. The reactor was sealed and heated to the indicated reaction temperature, then pressurized to the indicated PO_2 with oxygen gas. Agitation was initiated.

The selectivity (mole%) to N-phosphonomethylglycine was determined by dividing the moles of N-phosphonomethylglycine produced by the total moles of N-phosphonomethyliminodiacetic acid consumed and multiplying by 100. The yield (mole%) of N-phosphonomethylglycine was determined by dividing the moles of N-phosphonomethylglycine produced by the total moles of starting N-phosphonomethyliminodiacetic acid and multiplying by 100.

Examples 1 through 8

Examples 1 through 8, shown in Table 1, show the effect of varying the manganese catalyst concentration. In examples 1-4 the reaction temperature was 90 °C, the PO₂ was 100 psig (690 kPa), the initial N-phosphonomethyliminodiacetic acid concentration was 0.1M. The catalyst was a mixture of Mn(II) and Mn(III) acetate salts in a 1:1 mole ratio of Mn(II) and Mn(III). Examples 5-8 were run at the same conditions as 1-4, except that the PO₂ was 450 psig (3100 kPa) and the reaction temperature was 80 °C and the catalyst was Mn(II) acetate.

Table 1

Effect of Varying Catalyst Concentration				
Examples	Selectivity to N-phosphonomethylglycine (Mole%)	Manganese Concentration (M)	Initial Reaction Rate (Velocity, M/hr)	Yield of N-Phosphonomethylglycine (Mole %) at indicated time (h)
1	58	0.008	0.23	53(6)
2	82	0.004	0.10	75(6)
3	84	0.002	0.05	18(1½)
4	63	0.001	0.016	45(6)
5	83	0.02	0.30	83(2/3)
6	83	0.0067	0.10	81(½)
7	70	0.004	0.07	68(6)
8	74	0.002	0.034	68(6)

The data indicated that the reaction rate increases with the catalyst concentration. There appeared to be a first-order dependence of the reaction rate on the catalyst concentration.

Examples 9 through 13

Examples 9 through 13, shown in Table 2, illustrate the effect of initial pH on the reaction rate and selectivity to N-phosphonomethylglycine for a manganese catalyst. The reaction temperature was 80 °C, the PO₂ was 100 psig (690 kPa), the initial N-phosphonomethyliminodiacetic acid concentration was 0.1M, the reaction times are indicated and the manganese ion concentration was 0.004M. The mixture of manganese salts was the same as used in Example 1. The initial pH was adjusted using sodium hydroxide or sulfuric acid solutions. The data indicate that as the initial pH increases, the reaction rate increases, but the selectivity to N-phosphonomethylglycine decreases.

Table 2

Effect of Varying Initial pH				
Example	Initial pH	Initial Reaction Rate (M/h)	Yield of N-phosphonomethyl glycine (Mole %) at indicated time (h)	Selectivity to N-phosphonomethyl glycine (Mole %)(h)
9	1.20	0.0103	31(6)	49(6)
10	1.35	0.015	56(5)	66(5)
11	1.80	0.11	41(2½)	44(2½)
12	2.30	0.14	36(2½)	37(2½)
13	3.50	0.32	39(39)	41(½)

Examples 14 through 16

Examples 14 through 16, shown in Table 3, illustrate the effect of reaction temperature on reaction rates and selectivity to N-phosphonomethylglycine for a manganese catalyst. The PO_2 was 450 psig, the initial N-phosphonomethyliminodiacetic acid concentration was 0.1M and the manganese ion concentration was 0.067M. The form of the manganese salt was Mn(II)SO_4 and the pH was the unadjusted pH of the acid solution.

The data indicated that as the reaction temperature increased, the reaction rate increased.

Table 3

Effect of Varying Temperature				
Example	Temperature (°C)	Initial Reaction Rate (M/hr)	Selectivity to N-phosphonomethyl glycine (Mole %) at indicated time (h)	Yield of N-phosphonomethyl glycine (Mole %) at indicated time (h)
14	70	0.035	77 (5)	75(5)
15	80	0.093	83 (1½)	81(1½)
16	90	0.310	80 (½)	77(½)

Examples 17 through 22

Examples 17 through 22, shown in Table 4, illustrate the effect of PO_2 on selectivity to N-phosphonomethylglycine for a manganese catalyst. The reaction temperature was 80°C, the initial N-phosphonomethyliminodiacetic acid concentration was 0.1, the reaction time was as indicated which allowed for almost complete conversion of the N-phosphonomethyliminodiacetic acid, and the manganese ion concentration was 0.006 M. The form of the manganese salt was Mn(II)SO_4 and the pH was the unadjusted pH of the acid solution.

The data indicated that as the PO_2 increased, the selectivity to N-phosphonomethylglycine increased.

Table 4

Effect of Varying PO ₂			
Example	PO ₂ psig (kPa)	Selectivity to N-phosphonomethyl glycine (Mole %) at the indicated time (h)	Yield of N-phosphonomethyl glycine (Mole %) at the indicated time (h)
17	40(210)	56(6)	54(6)
18	70(450)	65(6)	63(6)
19	100(690)	68(6)	66(6)
20	130(890)	75(6)	73(6)
21	225(1550)	81(2)	78(2)
22	450(3100)	83(1½)	81(1½)

Examples 23 through 29 and Control 1

Examples 23 through 29 and Control 1, shown in Table 5, illustrate the effect of varying the form of the manganese catalyst on selectivity to N-phosphonomethylglycine. The reaction temperature was 90° C, the PO₂ was 100 psig (700 kPa); the initial concentration of N-phosphonomethyliminodiacetic acid was 0.1M, the manganese concentration was 0.004M and the reaction time was 1 h. The pHi was the unadjusted pH of the acid solution.

The Mn(III)chloro-(phthalocyaninato) (Control 1) was not catalytic.

Table 5

Effect of Varying Form of Manganese			
Example	Form	Selectivity to N-phosphonomethyl glycine (Mole %) at 1 h.	Selectivity at 6 h.
23	¹ Mn(II)/Mn(III)	43	75
24	Mn(II)acetate	18	75
25	Mn(III)acetate	20	75
26	Mn(II)sulfate	16	75
27	² Mn(II)(acac)	20	75
28	³ MnCl ₂ · 4H ₂ O	82	-
29	³ MnO ₂	70	73
Control 1	⁴ Mn(III)	1	<10

¹ Mn acetate, 50.50 mole ratio Mn(II)/Mn(III)

² Mn(II)bis(acetylacetonate)

³ PO₂ = 450 psig (3100 kPa) at 80° C and Mn concentration was 0.01M.

⁴ Mn(III)chloro-(phthalocyanato)

Examples 30 through 42

Examples 30 through 42, shown in Table 6, further illustrate the present invention. The initial pH, unless otherwise indicated, was the unadjusted pH at reaction temperature, the PO₂, unless otherwise indicated, is 100 psig (690 kPa), the initial concentration of N-phosphonomethyliminodiacetic acid was 0.1M, and the manganese catalyst was the mixture used in Example 1.

Table 6

Example	Run Time (h)	Catalyst Concentration(M)	Temperature (°C)	Yield (Mole%)	Conversion (Mole%)
30	1	.01	90	10	96
31	1	.02	80	42	97
32	1 ^a	.007	80	32	91
33	2	.01	70	8	95
34	2	.007	80	65	95
35	2 ^b	.007	70	74	96
36	2 ^c	.007	80	25	75
37	2 ^d	.007	80	22	63
38	2	.004	90	42	80
39	2	.002	90	60	75
40 ^e	2½	.007	80	85	100
41 ^f	1	.007	80	95	97
42 ^g	5	.07	80	19	84

a. pH_i = 2.3

b. PO₂ = 130 psig(810 kPa)

c. PO₂ = 40 psig(275 kPa)

d. pH_i = 1.35

e. PO₂ = 225 psig(1545 kPa)

f. PO₂ = 450 psig (3100 kPa)

g. Catalyst was Mn(II)acetylacetonate, the PO₂ was 450 psi(3000 kPa) and the initial concentration of N-phosphonomethyliminodiacetic acid was 0.5M.

Examples 43 through 65

Examples 43 through 65, shown in Table 7, illustrate the use of cobalt catalysts in the present invention. The initial concentration of N-phosphonomethyliminodiacetic acid was 0.1 M and the catalyst was Co(II)(SO₄). The pH was the unadjusted pH of the N-phosphonomethyliminodiacetic acid of the solution, unless otherwise indicated when it was adjusted with sodium hydroxide or sulfuric acid solution.

Table 7
Cobalt Catalysts

Example	Run Time (h)	Catalyst Concentration (M)	Temperature (°C)	Yield (Mole %)	Conversion (Mole %)	pH	PO ₂ (psi)
43	5.5	0.02	80	73	100	unadjusted	450
44	3.0	0.02	85	85	100	"	450
45	1.75	0.02	90	75	100	"	450
46	5.5	0.02	85	90	100	"	450
47	5	0.02	85	98	100	"	1000
48	2.0 ^a	0.02	85	21	31	"	450
49	5.5 ^b	0.02	85	74	98	"	300
50	3.0 ^c	0.036	90	87	100	"	450
51	4.0 ^c	0.048	80	64	97	"	450
52	5.0 ^d	0.125	85	52	99	"	450
53	18 ^f	0.5	100	16	100	6.25	100
54	18 ^e	0.5	100	28	98	1.80	100
55	18 ^e	0.5	100	16	100	2.25	100
56	18 ^e	0.5	100	0	100	4.00	100
57	18 ^e	0.5	100	35	98	1.09	100
58	18 ^e	0.5	100	9.9	22	0.77	100
59	18 ^e	0.5	100	17	98	1.7	100

Table 7 (cont'd.)
Cobalt Catalysts

Example	Run Time (h)	Catalyst Concentration (M)	Temperature (°C)	Yield (Mole %)	Conversion (Mole %)	pH	PO ₂ (psi)
60	18 ^c	0	100	0	98	9.00	100
61	18 ^e	0.01	100	20	40	0.44	100
62	2 ^f	0.01	100	28	98	1.80	100
63	28	0.01	100	26	98	1.80	100
64	18 ^h	0.01	100	26	98	1.74	100
65	5 ⁱ	0.2	85	66	99	1.7M	450

a The catalyst was Co(III)(acetylacetonate)₃.

b The initial N-phosphonomethyliminodiacetic acid concentration was 0.3M.

c The initial N-phosphonomethyliminodiacetic acid concentration was 0.4M.

d The initial N-phosphonomethyliminodiacetic acid concentration was 1.0M.

e The initial N-phosphonomethyliminodiacetic acid concentration was 0.5M, the catalyst was CoCl₂.

f The initial N-phosphonomethyliminodiacetic acid concentration was 0.5M and the catalyst was Co(NO₃)₂.

g The initial N-phosphonomethyliminodiacetic acid concentration was 0.5M and the catalyst was cobalt acetate.

h The initial N-phosphonomethyliminodiacetic acid concentration was 0.5M and the catalyst was CoBr₂.

i The initial N-phosphonomethyliminodiacetic acid concentration was 0.4M.

55 Examples 66 through 85

Examples 66 through 85, shown in Table 8, illustrate iron catalysts suitable for the present invention. The PO₂ was 100 psi (690 kPa), the catalyst concentration was 0.01 M, the reaction temperature was

100°C, the run time was 18 h, and the initial concentration of the N-phosphonomethyliminodiacetic acid was 0.5M, which formed a slurry. When NaBr was added, the concentration was also 0.01 M.

Table 8
Iron Catalysts

Example	Catalyst	Yield (mole %)	Conversion (mole %)	pH
66	Fe(SO ₄) ₂	21	36	6.25
67	"	18	28	10.0
68	"	6	14	5.0
69	" + NaBr	5	6	3.0
70	"	12	14	5.0
71	"	26	40	6.25
72	"	28	84	7.0
73	"	29	84	8.0
74	"	37	83	9.0
75	iron(III)(dicyano)bis (o-phenanthroline) tetrafluoroborate salt	6	12	6.25
76	"	8	10	7.0
77	"	3	12	9.0
78	"	3	12	10.0
79	K ₃ Fe(CN) ₆ ^a	3	14	3.0
80	"	8	24	5.0
81	"	21	46	6.3
82	"	30	76	7.0
83	"	37	80	9.0
84	"	32	80	10.0
85	Fe(SO ₄) ₂ +Al(NO ₃) ₃	21	72	6.0

^a Run time is 8 h.

Examples 86 through 106 and Control 2

Examples 86 through 106 and Control 2, shown in Table 9, illustrate nickel, chromium, ruthenium, aluminum, and molybdenum catalysts appropriate for the present invention. The conditions are as for those given in Table 8. The catalyst for Control 2, CuCl₂, appeared to be ineffective.

Table 9
Nickel Chromium, Ruthenium, Aluminum
and Molybdenum Catalysts

Examples	Catalyst	Yield (mole%)	Conversion (mole%)	pH
86	NiBr ₂	0.2	22	5.0
87	"	0.2	10	4.0
88	"	10	34	7.0
89	"	9	38	8.4
90	"	8	34	10.4
91	CrCl ₃	1	12	1.26
92	"	4	16	2.0
93	"	16	76	3.0
94	"	0.1	14	4.0
95	"	12	52	5.0
96	"	4	22	7.0
97	"	13	58	6.25
98	RuBr ₃	70	8	6.25
99	"	18	34	10.0
100	RuBr ₂ (Me ₂ SO) ₄	34	62	6.25
101	"	25	48	11.0
102	Al(NO ₃) ₃	11	34	6.25
103	" + NaCl	12	16	6.25
Control 2	CuCl ₂	0.2	14	6.25
104	K ₄ Mo(CN) ₈	4	22	4.0
105	"	32	48	6.0
106	"	10	30	9.0

Examples 107-109

Examples 107 through 109 shown in Table 10, illustrate vanadium catalysts suitable for the present invention. The reaction temperature was 70 °C, the PO₂ was 100 psi (690 kPa), the initial concentration of N-phosphonomethyliminodiacetic acid was 0.5 M, the catalyst concentration was 0.033M.

Table 10

Vanadium Catalysts				
Examples	Catalyst	Run Time (h)	Yield (mole%)	Conversion (mole%)
107	VO(acetylacetonate) ₂	2	40	67
108	VOSO ₄ (hydrate)	2.25	42	94
109	VOSO ₄ (hydrate) ^a	5	54	91

^a The initial concentration of N-phosphonomethyliminodiacetic acid was 0.15 M and the concentration of catalyst was 0.015M.

Examples 110 and 111

Examples 110 and 111 shown in Table 11 illustrate cerium catalysts suitable for the present invention. The reaction temperature was 90 °C and the PO₂ was 130 psi (897 kPa).

Table 11

Cerium Catalysts

Example	Catalyst	Run Time (h)	Catalyst Concentration (M)	N-phosphonomethyl- iminodiacetic acid Concentration (M)	Yield (mole %)	Conversion (mole %)
110	$\text{Ce}(\text{NH}_4)_4(\text{SO}_4)_4$	3	0.1	1.0	7	45
111	$\text{Ce}(\text{NH}_4)_4(\text{SO}_4)_4$	3	0.01	0.1	30	80

55 Claims

1. A process for the production of N-phosphonomethylglycine comprising contacting N-phosphonomethyliminodiacetic acid with molecular oxygen in the presence of a catalyst selected from

salts and salt complexes of manganese, cobalt, iron, nickel, chromium, ruthenium, aluminum, molybdenum, vanadium and cerium, wherein the reaction temperature is in the range of about 25° to 150° C, the reaction pressure is in the range of about atmospheric (101 kPa) to about 20,700 kPa (3000 psig), and the partial pressure of oxygen is in the range of about 0.1 to 7.0.

2. A process of Claim 1 wherein the catalyst is at least one of a manganese (II), a manganese (III), a vanadium (IV), a vanadium (V), a cerium (III) or a cerium (IV) salt.
3. A process of Claim 2 wherein the catalyst is a mixture of manganese (II) and manganese (III) salts.
4. A process of Claim 2 wherein the catalyst is selected from manganese acetate, manganese sulfate, manganese (II) bis (acetylacetonate) and manganese chloride.
5. A process of Claim 1 wherein the catalyst is a cobalt catalyst.
6. A process of Claim 5 wherein the catalyst is selected from cobalt chloride, cobalt bromide, cobalt sulfate, cobalt bis(acetylacetonate) and cobalt acetate.
7. A process of Claim 1 wherein the catalyst is selected from vanadiumoxy (acetylacetonate) and vanadiumoxysulfate (hydrate).

Patentansprüche

1. Verfahren zur Herstellung von N-Phosphonomethylglycin umfassend das Inberührungbringen von N-Phosphonomethyliminodiessigsäure mit molekularem Sauerstoff in Anwesenheit eines Katalysators ausgewählt aus Salzen und Salzkomplexen von Mangan, Kobalt, Eisen, Nickel, Chrom, Ruthenium, Aluminium Molybdän, Vanadium und Cer, wobei die Reaktionstemperatur im Bereich von etwa 25 bis 150° C liegt, der Reaktionsdruck im Bereich von etwa atmosphärisch (101 kPa) bis etwa 20.700 kPa (3000 psig) ist und der Sauerstoffpartialdruck im Bereich von etwa 0,1 bis 7,0 liegt.
2. Verfahren nach Anspruch 1, in dem der Katalysator zumindest eines von Mangan(II)-, Mangan(III)-, Vanadium(IV)-, Vanadium(V)-, Cer(III)- oder Cer(IV)-Salz ist.
3. Verfahren nach Anspruch 2, in dem der Katalysator eine Mischung von Mangan(II)- und Mangan(III)-salzen ist.
4. Verfahren nach Anspruch 2, in dem der Katalysator ausgewählt ist aus Manganacetat, Mangansulfat, Mangan(II)bis(acetylacetonat) und Manganchlorid.
5. Verfahren nach Anspruch 1, in dem der Katalysator ein Kobaltkatalysator ist.
6. Verfahren nach Anspruch 5, in dem der Katalysator ausgewählt ist aus Kobaltchlorid, Kobaltbromid, Kobaltsulfat, Kobalt-bis(acetylacetonat) und Kobaltacetat.
7. Verfahren nach Anspruch 1, in dem der Katalysator ausgewählt ist aus Vanadiumoxy(acetylacetonat) und Vanadiumoxysulfat(hydrat).

Revendications

1. Procédé pour la production de N-phosphonométhylglycine comprenant la mise en contact d'acide N-phosphonométhyliminodiacétique avec de l'oxygène moléculaire en présence d'un catalyseur choisi parmi des sels et des complexes sels de manganèse, de cobalt, de fer, de nickel, de chrome, de ruthénium, d'aluminium, de molybdène, de vanadium et de cérium; dans lequel la température de réaction est comprise entre environ 25° et 150° C, la pression de réaction est comprise entre environ la pression atmosphérique (101 kPa) et environ 20.700 kPa (3000 psig) et la pression partielle de l'oxygène est comprise entre environ 0,1 et 7,0.
2. Procédé selon la revendication 1, dans lequel le catalyseur est au moins un des éléments constitué par

un sel de manganèse (II), un sel de manganèse (III), un sel de vanadium (IV), un sel de vanadium (V), un sel de cérium (III) ou un sel de cérium (IV).

- 5 3. Procédé selon la revendication 2, dans lequel le catalyseur est un mélange de sels de manganèse (II) et de manganèse (III).
4. Procédé selon la revendication 2, dans lequel le catalyseur est choisi parmi l'acétate de manganèse, le sulfate de manganèse, le bis (acétylacétonate) de manganèse (II) et le chlorure de manganèse.
- 10 5. Procédé selon la revendication 1, dans lequel le catalyseur est un catalyseur de cobalt.
6. Procédé selon la revendication 5, dans lequel le catalyseur est choisi parmi le chlorure de cobalt, le bromure de cobalt, le sulfate de cobalt, le bis(acétylacétonate) de cobalt et l'acétate de cobalt.
- 15 7. Procédé selon la revendication 1, dans lequel le catalyseur est choisi parmi l'oxy(acétylacétonate) de vanadium et l'oxysulfate de vanadium (hydrate).

20

25

30

35

40

45

50

55

THIS PAGE BLANK (USPTO)